



## Use of First Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP)

### APP Measure Description<sup>1</sup>

Assesses the percentage of children and adolescents newly started on antipsychotic medications without a clinical indication who had documentation of psychosocial care as first-line treatment.

### Why is APP Important?<sup>1</sup>

Although antipsychotic medications may serve as effective treatment for a narrowly defined set of psychiatric disorders in children and adolescents, they are often prescribed for nonpsychotic conditions for which psychosocial interventions are considered first-line treatment.

Safer, first-line psychosocial interventions may be underutilized, and children and adolescents may unnecessarily incur the risks associated with antipsychotic medications.

### Best Practices

- ✓ When prescribed, antipsychotic medications should be part of a comprehensive, multi-modal plan for coordinated treatment that includes psychosocial care.
- ✓ Psychosocial care, which includes behavioral interventions, psychological therapies and skills training, among others, is the recommended first-line treatment option for children and adolescents diagnosed with nonpsychotic conditions such as attention-deficit disorder and disruptive behaviors.
- ✓ Periodically review the ongoing need for continued therapy with antipsychotic medications.
- ✓ Assess the need for Case Management and refer if necessary.
- ✓ Ensure progress notes are completed and accurate.

All summaries of the measure contained herein are reproduced with permission from HEDIS Volume 2: Technical Specifications for Health Plans by the National Committee for Quality Assurance (NCQA). HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA)

<sup>1</sup> Source: [ncqa.org/hedis/measures](https://www.ncqa.org/hedis/measures)

<sup>2</sup> Source: HEDIS MY 2024 Tech Specs Manual Vol. 2

## Numerator Compliance<sup>2</sup>

Psychosocial care (Psychosocial Care Value Set) or residential behavioral health treatment (Residential Behavioral Health Treatment Value Set) in the 121-day period from 90 days prior to the IPSD through 30 days after the IPSD.

Index prescription start date (IPSD):  
The earliest prescription dispensing date for an antipsychotic medication where the date is in the intake period and there is a negative medication history.

## Numerator Codes<sup>2</sup>

Refer to HEDIS Numerator Codes Dashboard on the Trillium HEDIS Resources website.

## Data Collection Method<sup>2</sup>

Administrative (Claims)



## Trillium Percentages/NCQA National Averages<sup>1</sup>

APP	Calendar Year	Trillium	NCQA National Average
Use of First-line Psychosocial Care for Children and Adolescents in Antipsychotics	2022	54.3	57.3
	2021	48.8	58.6

All summaries of the measure contained herein are reproduced with permission from HEDIS Volume 2: Technical Specifications for Health Plans by the National Committee for Quality Assurance (NCQA). HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA)

<sup>1</sup> Source: [ncqa.org/hedis/measures](https://www.ncqa.org/hedis/measures)

<sup>2</sup> Source: HEDIS MY 2024 Tech Specs Manual Vol. 2